Report card
on the WHO
Framework
Convention
on Tobacco
Control



Introduction

Tobacco use is the single most preventable cause of death in the world today, and is estimated to kill more than five million people every year – more than tuberculosis, HIV/AIDS and malaria combined. By 2030, the death toll will exceed eight million a year, unless urgent action is taken. Implementation of the World Health Organization Framework Convention on Tobacco Control (WHO FCTC) can reverse this devastating epidemic if it becomes a top public health priority especially for countries' political leaders.

The Senegal Report Card on the WHO FCTC provides descriptive analysis of country-level data that reflect the status and progress on some key articles of the WHO FCTC. This is one of a series of Report Cards for countries in the WHO African Region.

\frown		en	4
	mt	en	TQ
	m	$\mathbf{c}_{\mathbf{II}}$	LO

1
2
3
4
5
6
7
8
9
10

Senegal entry into force of the WHO FCTC



27 April 2005

WHO Framework Convention on Tobacco Control

Compliance dates for Senegal

27 April 2008 Article 11: Health warnings on tobacco products and packaging

27 April 2010 Article 13: Bans on tobacco advertising, promotion & sponsorship

27 April 2010 Article 21: Next Report to the Conference of the Parties

Background

1. WHO Framework

Convention on Tobacco Control (WHO FCTC)

The WHO FCTC is an evidence-based treaty that was developed in response to the globalization of the tobacco epidemic. It was adopted by the World Health Assembly in 2003 and became international law on 27 February 2005. The treaty contains articles aimed at both reducing the supply of and demand for tobacco. This Report describes country progress on the following key WHO FCTC Articles and relevant Guidelines:

Article 5.3: Protect public health policies from tobacco industry interference;

Article 6: Price and tax measures to reduce the demand for tobacco;

Article 8 provisions and Guidelines: Protection from exposure to tobacco smoke;

Article 11 and Guidelines: Packaging and labelling of tobacco products;

Article 13 and Guidelines: Tobacco advertising, promotion and sponsorship;

Article 20: Research, surveillance and exchange of information.

2. Reporting Instrument of the WHO FCTC

The WHO FCTC requires countries that are Parties to the Convention (Parties) to submit periodic reports to the Conference of the Parties (COP). The objective of these reports is to "enable Parties to understand and learn from one another's experiences in implementing the WHO FCTC."

In accordance with WHO FCTC Article 21, Parties report initially 2 years after entry into force of the WHO FCTC for that Party, and then every 3 years, i.e.:

Phase 1 (Group 1 questions): within 2 years of entry into force;

Phase 2 (Group 2 questions): within 5 years of entry into force;

Phase 3 (Group 3 questions): within 8 years of entry into force.

Reporting tools are provided at the WHO FCTC COP website: www.who.int/fctc/reporting/reporting_instrument/en/index.html

ACRO WHO Regional Office for Africa

CDC United States Centers for Disease Control and Prevention

COP WHO FCTC Conference of the Parties

FCTC [WHO] Framework Convention on Tobacco Control

GATS Global Adult Tobacco Survey

GHPS Global Health Professional Students Survey

GSPS Global School Personnel Survey
GTSS Global Tobacco Surveillance System
GYTS Global Youth Tobacco Survey

GTCR WHO Report on the Global Tobacco Epidemic

NRT Nicotine replacement therapy
TFI WHO Tobacco Free Initiative
WHO The World Health Organization

Definitions

Ad valorem tax: Tax applied on the value of items (i.e. a percentage of the price)

Current smoker: Anyone who currently smokes any tobacco product on some or all days

Public place: A place which the public is entitled to use or which is open to, or used by the public or a section of the public

Public place, indoor: Public places that are covered by a roof and one or more walls/sides

Public transport: Any vehicle used at any time by members of the public

Second-hand smoke: Both smoke from the burning end of a cigarette or other tobacco product and smoke exhaled by the smoker

Smoke-free air: Air that is 100% smoke-free and that cannot be sensed or measured

Specific tax: Tax applied on unit quantities of items (e.g. \$1 per pack of 20 cigarettes)

Tobacco advertising & promotion: Commercial communication, recommendation or action with the aim, effect or likely effect of promoting a tobacco product or tobacco use directly or indirectly

Tobacco industry: Manufacturers, wholesale distributors and importers of tobacco products

Tobacco sponsorship: Contribution to any event, activity or individual with the aim, effect or likely effect of promoting a tobacco product or tobacco use directly or indirectly

Workplace: Place used by people during their employment or work, including vehicles

3. Global Tobacco Surveillance System (GTSS)

WHO, the US Centers for Disease Control and Prevention (CDC), and others began development of the GTSS in 1999 with the purpose of enhancing the capacity of countries to design, implement, and evaluate their national comprehensive tobacco control action plan and to monitor some key articles of the WHO FCTC. The GTSS includes four surveys:

Global Youth Tobacco Survey (GYTS): school-based survey of students aged 13–15 years that provides information on demographics, tobacco use prevalence, and knowledge and attitudes.

Global School Personnel Survey (GSPS): provides demographics, tobacco use prevalence, knowledge and attitudes information, as well as information on school policy and curriculum.

Global Health Professional Students Survey (GHPS): school-based survey of 3rd-year students pursuing advanced degrees in dentistry, medicine, nursing, or pharmacy.

Global Adult Tobacco Survey (GATS): household survey that tracks tobacco use prevalence, exposure to risk, second-hand smoke, cessation, risk perceptions, knowledge and attitudes, exposure to media and price as well as taxation issues.

4. WHO Reports on the Global Tobacco Epidemic, 2008 and 2009

These WHO reports detail information on internationally comparable prevalence estimates and data that allows assessment of country compliance with policies on smoke-free environments, treatment of tobacco dependence, health warnings and packaging, advertising, promotion and sponsorship bans, price and taxation levels, and key national capacity indices.

Status of WHO FCTC in Senegal

Article	Requirement	Senegal status
Article 5.1 National strategies, plans, programmes and coordinating mechanism	Each Party shall develop, implement, periodically update and review comprehensive multisectoral national tobacco control strategies, plans and programmes, establish or reinforce and finance a national coordinating mechanism or focal points for tobacco control.	National agency with staff as well as national strategies, plans and programmes on tobacco control.
Article 6 Price and tax measures to reduce demand for tobacco products	Take account of national health objectives concerning tobacco control and adopt or maintain measures which may include implementing tax and price policies on tobacco products so as to contribute to the health objectives aimed at reducing tobacco consumption; and prohibiting or restricting taxand duty-free tobacco products.	Total tax on the most sold brand is 28 %.
Article 8 Protection from exposure to tobacco smoke	Adopt and implement measures, providing for protection from exposure to tobacco smoke in indoor workplaces, public transport, indoor public places and, as appropriate, other public places.	Smoke-free policies in health-care facilities.
Article 11 Packaging and labelling of tobacco products	Adopt measures within 3 years of entry into force that require display of rotated series of health warnings and other messages on tobacco product packaging that cover at least 30% of the principle display areasideally 50% or more, and include pictures or pictogramsand that prevent false, misleading or deceptive packaging and labelling.	
Article 13 Tobacco advertising, promotion and sponsorship	Undertake a comprehensive ban of all tobacco advertising, promotion and sponsorship within 5 years of entry into force.	Ban on free distribution of tobacco products.

Senegal National action is critical in order to achieve the vision embodied in the WHO FCTC.	National strategies, plans and programmes on tobacco control	National agency on tobacco control	National budget for tobacco control
	Senegal has established national strategies, plans and programmes on tobacco control.	Senegal has a national agency with eight full-time equivalent employees.	In 2009, the budget was US \$77 201 (at official exchange rate).
			\$

National action

Building national capacity to carry out effective and sustainable national tobacco control programmes is one of the most significant measures required to combat the tobacco epidemic. Successful implementation of the WHO FCTC requires establishing national a coordinating mechanism with an official government mandate for developing and coordinating a plan of as well for action as building national a infrastructure to implement and evaluate the plan.

Cessation

The Senegal population a toll-free has access to quitline. **Nicotine** replacement therapy, bupropion and vareniclin are legally sold in the country. Smoking cessation support is available in some primary hospitals, care facilities, offices of health professionals and community centres.

In 2002, 87.2% of the currently smoking students expressed a desire to stop. This proportion decreased to 77.4% in 2007.

Senegal has a national toll-free quitline.


Tobacco use prevalence


Article 20 WHO FCTC on research, surveillance and exchange of information requires Parties to establish, as appropriate, programmes for national, regional and global surveillance of the magnitude, patterns, determinants and consequences of tobacco consumption and exposure to tobacco smoke.

AFRO recommends that countries obtain nationally representative and population-based periodic data on key indicators of tobacco use for youth and adults. AFRO also recommends that countries continue to successfully participate in the GYTS, expand participation in the GSPS, GHPS and GATS as well as other representative national surveys. Countries are also encouraged to report the findings widely and utilize them to inform tobacco control policy development, programme planning and evaluation.

Adult (2008)

Youth (2002/2007)





Adult tobacco use

Among adults of 18 years and above, prevalence of current tobacco smoking was 11.6% in 2003 with 22.2% males and 1.7% females.

Youth tobacco use

Between 2002 and 2007, 13-15 years old students reported a decrease in prevalence of current cigarette smoking but an increase in prevalence of current use of other tobacco products. In 2002, 13.2% of students (boys=20.2%; girls=4.4%) currently smoked cigarettes and 5.4% (boys=7.3%; girls=2.9%) currently used other tobacco products.

In 2007, 7.5% of students (boys=12.1%; girls=2.7%) currently smoked cigarettes and 9.3% (boys=11.7%; girls=7.7%) currently used other tobacco products.

Cigarette smoking among Senegal youth decreased, but use of other tobacco products increased between 2002 and 2007.

Smoke-free policies

Countries should protect people from tobacco smoke

- Every person has a right to breathe air free of tobacco smoke.
- In addition to protecting the health of non-smokers, smoke-free environments encourage smokers to quit.
- Evidence from pioneering countries shows that smoke-free laws do not harm businesses and are popular with the public.
- Permitting smoking in designated areas undermines the benefit of smoke-free environments



Senegal law

Article 8 of the WHO FCTC mandates protection from exposure to tobacco smoke in indoor workplaces, public transport and indoor and other public places. Guidelines for this Article were unanimously adopted by the Conference of the Parties in July 2007.

AFRO recommends that countries enact and enforce completely smoke-free environments in all indoor public places including workplaces, restaurants and bars. AFRO further recommends that all countries should become compliant with the requirements of Article 8 guidelines, and that 100% smoke-free environments should become the norm in all societies.

Senegal has national laws that completely ban smoking indoors in health-care facilities. Educational facilities, universities, governmental facilities, indoor offices, other indoor workplaces, restaurants, pubs & bars are not covered by this regulation. There are subnational jurisdictions that have authority to adopt and implement laws that ban tobacco smoking in such places.

Youth exposure to second-hand smoke

Both in 2002 and 2007, students reported high exposure to second-hand tobacco smoke.

In 2002, close to half (45.8%) of students lived in homes where others smoked in their presence and over 6 in 10 students (62.6%) were exposed to smoke outside their homes. Similarly, in 2007, almost half of the students (47.6%) lived in home where others smoked in their presence and nearly half of the students (48.3%) were exposed to smoke outside of their homes. In 2002, 89.5% of students thought smoking should be banned from public places. This decreased to 85.9% in 2007.

Health warnings and education

should Countries about the warn dangers of tobacco use and exposure to tobacco smoke



Mauritius health warning

- ONE specific warning with large, clear
- NO mandate for principal display surface
- NO pictures or pictograms
- NO ban on false, misleading or deceptive packaging and labeling
- Rotating, specific warnings and large, clear
- 65% of display surface
- Pictures or pictograms
- Ban false, misleading or deceptive packaging and labeling

Best practice

Article 11 WHO FCTC requires Parties to implement clear health warnings and measures that remove false or misleading promotion of tobacco products.

Relatively few tobacco users fully grasp the health dangers. Hardhitting anti-tobacco advertising and graphic pack warnings reduce the number of children who begin smoking and increase the number of smokers who quit.

Pictures are more powerful deterrents than words on tobacco packaging warnings, but globally, only 15 countries mandate pictorial warnings. Just five countries meet the highest standards for pack warnings.

Countries should also promote and strengthen public awareness of tobacco control issues, using all available communication tools.

Senegal labelling

There is no ban on use of deceitful terms, such as 'low tar', 'light' or 'mild' on tobacco product packaging in Senegal. The mandates one specific warning tobacco product on packages. The warning appears on each package and any outside packaging and labelling used in retail sale and is written in the principal language of the country. The country does not mandate that a certain percentage of packaging principal display area be covered by a health warning or that warning describe the harmful effects of tobacco use or graphic warnings or rotation of warnings.

Senegal education

In 2002, 48.4% and 41.6% of students had been taught in school about dangers of smoking and effects of tobacco use, respectively. In 2007, 55.3% and 42.7% of students had been taught in school about dangers of smoking and effects of tobacco use, respectively.



Bans in Senegal

Senegal does not have a national ban on direct tobacco advertising. There is a ban on free distribution of tobacco products in mail or by other means. There is no ban on other forms of promotion tobacco and sponsorship.

Youth exposure

Exposure of 13-15 years old students to most tobacco advertising increased between 2002 and 2007. In 2002, 72.4% of students had seen pro-cigarette advertising on billboards and 57.1% in newspapers or magazines.

Similarly, in 2007, 75% of students had seen procigarette advertising on billboards and 63.3% in newspapers or magazines. In 2002, 23.3% of students reported having an object with a cigarette brand logo, but by 2007 this proportion had decreased to 18.9%.

Advertising, promotion & sponsorship

Countries should enforce comprehensive bans on tobacco advertising, promotion and sponsorship

- Widespread advertising falsely associates tobacco with desirable qualities.
- Studies have found that advertising bans can lower tobacco consumption.
- About half the children of the world live in countries that do not ban free distribution of tobacco products.



Best practices

Tobacco taxes are the most effective way to reduce tobacco use, especially among young people and the poor.

Tobacco tax increases also increase government revenues. Only four countries, representing 2% of the world's population, have tax rates greater than 75% of the retail price.

In countries with available information, tobacco tax revenues are more than 500 times higher than spending on tobacco control. In low- and middle-income countries, tobacco tax revenues are more than 9000 and 4000 times higher than spending on tobacco control, respectively.

Senegal taxes

In 2009 the price of the most sold brand of cigarettes was about US \$1.27. The total taxes on most sold brand are 28%; total excise on most sold brand is 13%.

Increasing tobacco taxes by 10% generally decreases tobacco consumption by about 8% in low- and middleincome countries.

^{*20-}cigarette pack of most sold brand

Senegal policy recommendations

National action

AFRO recommends that successful implementation of the WHO FCTC requires establishing a national coordinating mechanism with an official government mandate for developing and coordinating a plan of action as well as for building a national infrastructure to implement and evaluate the plan.

Senegal should periodically update and review their national strategies, plans and programmes.

Smoke-free policies

AFRO recommends that countries become compliant with WHO FCTC Article 8 requirements and its guidelines and enact and enforce completely smokefree environments in all indoor public places including workplaces, restaurants and bars.

Senegal should fully comply with WHO FCTC Article 8 requirements and its guidelines.

Health warnings and education

AFRO recommends that countries fully comply with WHO FCTC Article 11 requirements and its guidelines and implement large, prominent, strong and effective graphic health warnings on packages, require effective package warning labels, implement countertobacco advertising, and obtain free media coverage of anti-tobacco activities.

Senegal should fully comply with WHO FCTC Article 11 requirements and its guidelines.

Bans on advertising, promotion and sponsorship

AFRO recommends that countries enact and enforce effective legislation that comprehensively bans any form of direct tobacco advertising, promotion and sponsorship; also, enact and enforce effective legislation to ban indirect tobacco advertising, promotion and sponsorship in line with WHO FCTC Article 13.

Senegal should fully comply with WHO FCTC Article 13 requirements and its guidelines.

Tobacco prices and taxes

AFRO recommends that countries increase tax rates for tobacco products to be at least 70% of retail prices and ensure that they are adjusted periodically to keep pace with inflation and rise faster than consumer purchasing power. It is also recommended that countries strengthen tax administration to reduce the illicit trade in tobacco products.

In addition, countries should protect public health policies from tobacco industry interference.

Senegal should raise tobacco taxes, ensure tobacco taxes are adjusted periodically, strengthen tax administration and protect public health policies from tobacco industry interference.

References

- Global Youth Tobacco Survey (GYTS) 2002 Fact Sheet: Senegal (Ages 13-15)
- Global Youth Tobacco Survey (GYTS) 2007: Senegal (Ages 13-15)
- MPOWER: A policy package to reverse the tobacco epidemic, WHO 2008
- WHO report on the global tobacco epidemic, 2008 - The MPOWER package, WHO 2008
- WHO report on the global tobacco epidemic, 2009: implementing smoke-free environments, WHO 2009
- WHO Framework Convention on Tobacco Control, WHO 2003
- Parties' reports on WHO FCTC implementation: http://www.who.int/fctc/reporting/party-report/en/ (accessed July 2010)