


ADOLESCENT HEALTH IN UR TANZANIA 2018



Country profile: An adolescent is a person between 10-19 years of age

T-1-1	4.4400
Total surface area (million sq. km)	1.1109
Total population (000's)	55,572.2010
Population density (persons/sq km)	60.8263
Urbanized population (%)	32.3160
Adolescent population (000's)	12,504.9190
Adolescent as % of total population	22.5021
Young people (10-24 years) (000's)	17,008.2360
Young people as % of total population	31.8101
Annual average growth rate of population (%)	2.9256
Crude birth rate (births per 1,000 population)	38.1860
Life expectancy at birth (years)	65.6750
Median age of the total population (years)	17.2910
Birth rate per 1000 girls 15-19 years olds	72.1000





Young people as % of total population




for injury prevention in adolescents

Early marriage








Median age at 1st sexual intercourse (years)

Percentage of 15-19 year olds M currently married or in union (%) 23 2


Pregnancy

Percentage of 15-19 year olds who have given birth/who are pregnant with their first child



Girls of 15-19 year olds who are pregnant with their first child Women girls of 15-19 years who have given birth

Contraception



HIV/STIs

	Compr	Comprehensive knowledge of HIV			
	15-19 years		20-24	years	
	Female	Male	Female	Male	
2007	35.8	41.3	42.6	47.2	
2010	46.3	40.9	50.4	45.4	
2003	32.4	35.2	41.0	44.1	
2012	36.8	41.9	44.2	53.0	
	Condom use among sexually active during premarital sex (%)				

	during premarital sex (%)				
	15-19 years		20-24	Years	
	Female	Male	Female	Male	
2007	49.4	40.3	47.4	56.6	
2010	50.2	46.1	47.6	60.4	
2012	60.2	50.7	53.8	65.4	
2016	37.1	3/1/6	35.8	19.5	

Prevalence (%) of HIV in 15-19 year olds



Prevalence (%) of STIs among 15-19 year



Unmet need for contraception





ADOLESCENT HEALTH IN UR TANZANIA 2018



Malnutrition

		15-19	years	20-24	years
		Female	Male	Female	Male
	1996	8.6		9.5	
Prevalence (%) of chronically malnourished (thin, BMI<18.5)	2010	17.6		9.3	
	2016	18.0		8.0	
	1996	5.2		10.0	
Prevalence (%) of overweight (BMI between 25.0-29.9)	2010	8.0		15.2	
	2016	9.1		15.4	
Prevalence (%) of iron deficiency anemia (Hqb<12.0 q/dl for non-pregnant women and	2010	42.2		40.5	
Hgb<11.0 g/dl for pregnant women)	2016	47.3		46.3	

Substance use

			Female	Male
Total alcohol consumption among 15-19 year olds (litres of pure alcohol)	15-19 years	2010	3.3	9.0
	15-19 years	2010	0.3	2.0
Prevalence cigarette or any		2015	0.3	
other tobacco product use (%)	20-24 years	2010	0.4	14.5
		2015	0.6	

Violence



Percentage of married adolescents who agree that husband is justified in beating his wife





59% 50%

			Female
% of married women who have ever experienced physical or sexual or emotional violence committed by their husband or partner	15-19 years	2010	43.8
		2015	44.2
	20-24 years	2010	46.0
		2015	49.1
Prevalence of victims women of femal genital mutilation (%)	15-19 years	2010	7.1
		2016	4.7
	20-24 years	2010	11.0
		2016	7.3

Injuries and trafic accidents

Adolescent household members injured and unable to carry out normal daily activities of >6 months (%)	
Adolescent household members injured or killed in accidental fall (%)	
Adolescent household members injured or killed in road traffic accident (%)	

Immunization

Proportion of adolescent girls vaccinated for HPV to



Mental health

	Female	Male
Prevalence of childhood mental illness among 15-19 year olds (%)		
Felt seriously worthless/hopeless or wishes to be dead among15-19 year olds (%)		



Socioeconomic determinants of nealth				
		Female	Male	
Literacy rate among 15-19 year olds (%)	2016	84.0	81.6	
Percent 15-19 year currently employed (%)	2016	44.9	64.0	
Information media use at least once/week among adolescents (%)	2016	56.7	61.4	
Social security health insurance among 15-19 year olds (%)	2015	1.2	1.8	

Trend in Mo	rtality rates per	100,000 in 15-	19 year olds
1996	2004	2010	2016
2.4 2.3	2.5 1 :7	3.5	1.3 1.2

Leading causes of girls' death

Rank cause name	number of deaths an	d death rate age	10-19
marin, caase marrie,	manner or acaths and	a acath rate, age	10 13

Rank	Cause name	Number of deaths	Death rate/100, 000 pop
1	Lower respiratory infections	1,216	19.4
2	Malaria	1,140	18.2
3	Diarrhoeal diseases	1,105	17.7
4	HIV/AIDS	1,077	17.2
5	Maternal conditions	779	12.4

Leading causes of boys' death

Rank, cause name, number of deaths and death rate, age 10-19			
Rank	Cause name	Number of deaths	Death rate/100, 000 pop
1	Diarrhoeal diseases	1,421	22.8
2	Lower respiratory infections	1,224	19.6
3	HIV/AIDS	1,187	19.0
4	Malaria	961	15.4
5	Road injury	894	14.3



For further information: